

An indirect limit on the amplitude of primordial Gravitational Wave Background from CMB-Galaxy Cross Correlation

Asantha Cooray,^{1,*} Pier-Stefano Corasaniti,^{2,†} Tommaso Giannantonio,^{3,‡} and Alessandro Melchiorri^{4,§}

¹*Dept. of Physics & Astronomy University of California, Irvine, CA 92697-4575*

²*ISCAP, Columbia University, 550 West 120th Street, New York, NY, 10027 (USA)*

³*Dipartimento di Fisica “G. Marconi”, Università di Roma “La Sapienza”, Ple Aldo Moro 5, 00185, Rome, Italy.*

⁴*Dipartimento di Fisica “G. Marconi” and INFN, sezione di Roma, Università di Roma “La Sapienza”, Ple Aldo Moro 5, 00185, Roma, Italy.*

While large scale cosmic microwave background (CMB) anisotropies involve a combination of the scalar and tensor fluctuations, the scalar amplitude can be independently determined through the CMB-galaxy cross-correlation. Using recently measured cross-correlation amplitudes, arising from the cross-correlation between galaxies and the Integrated Sachs Wolfe effect in CMB anisotropies, we obtain a constraint $r < 0.5$ at 68 % confidence level on the tensor-to-scalar fluctuation amplitude ratio. The data also allow us to exclude gravity waves at a level of a few percent, relative to the density field, in a low - Lambda dominated universe ($\Omega_\Lambda \sim 0.5$). In future, joining cross-correlation ISW measurements, which captures cosmological parameter information, with independent determinations of the matter density and CMB anisotropy power spectrum, may constrain the tensor-to-scalar ratio to a level above 0.05. This value is the ultimate limit on tensor-to-scalar ratio from temperature anisotropy maps when all other cosmological parameters except for the tensor amplitude are known and the combination with CMB-galaxy correlation allows this limit to be reached easily by accounting for degeneracies in certain cosmological parameters.

PACS numbers:

I. INTRODUCTION

One of the major predictions of inflation is the existence of a stochastic background of gravitational waves (GW) (see e.g. [1], [2]). The amplitude of these tensor fluctuations is proportional to the square of the energy scale of inflation (see e.g. [3]). Furthermore, the ‘tilt’ of the GW spectrum (and of the scalar as well) can give us direct information on the slope of the inflaton potential up to second derivatives. Therefore the detection of this stochastic background would provide valuable information on the inflationary scenario shedding light on the physics at $\sim 10^{16}$ GeV (see e.g. [4]).

The stochastic GW background leaves a signature on the CMB through different mechanisms. In principle a clear detection can be obtained from measurements of the B-polarization. Tensor perturbations also induce temperature fluctuations through a Sachs-Wolfe effect and an integrated one (for a review see [5]). The former is localized by the visibility function at the last scattering surface and consequently depends on the recombination history. Since it occurs in a relatively short period of time the amplitude of this effect is negligible at all scales. On the contrary GW produce most of the signal through the integrated effect which is sensitive to the evolution of gravitational waves from the time of last scattering to

today. Since the observed temperature anisotropy power spectrum is the sum of both the scalar and tensor components, for relatively low amplitudes of the tensor modes the scalar fluctuations dominate the spectrum and the tensor contribution is dwarfed. Similarly, if there is a non-negligible contribution from GWs, the sum of scalar and tensor spectra would lower the predicted amplitude of the acoustic peaks relative to the angular scales of few or more degrees. Thus, the ratio of tens of degree to sub-degree fluctuations can be used to constrain the presence of GW.

With the advent of WMAP, several studies have addressed whether there is a significant contribution of GWs in the CMB spectra (see e.g. [6, 7, 8, 9]). Despite the high quality of these data, it still remains difficult to obtain a robust constrain on the amplitude of the tensor perturbations due to the degeneracies with other cosmological parameters. For instance, both the scalar spectral index n_S and its running are strongly degenerate with the amplitude of the GW background. In [10], combining the CMB data with the 2dF matter power spectrum yield a 2σ constraint on the *tensor – to – scalar* ratio $r < 0.9$ [39]. The combination of WMAP data with Sloan Ly- α forest, lensing bias, and galaxy clustering data constrain $r < 0.36$ at the 95% confidence level in a flat cosmological model [11]. Measurements of B-mode CMB polarization from future experiments such as the *Planck* satellite, or in the long-term the CMBpol mission as part of NASA’s *Beyond Einstein* Program, will provide either a detection of the tensor modes or the best constrain on their amplitude [12]. Concept studies are already under way to consider the possibility for a direct detection of the relic gravitational wave background present today with

*Electronic address: acooray@uci.edu

†Electronic address: pierste@phys.columbia.edu

‡Electronic address: tommaso.giannantonio@roma1.infn.it

§Electronic address: alessandro.melchiorri@roma1.infn.it

space-based Laser interferometers as part of a second-generation mission to followup NASA's Laser Interferometer Space Antenna (LISA; [13]). In the meantime, it is worthwhile to investigate methods that could independently bound the GW contribution and/or break some of the degeneracies which affect present constraints. This would also be useful in planning observational strategies of current and upcoming CMB experiments.

The scalar and tensor temperature anisotropy power spectra are characterized by a nearly constant plateau on the large angular scales. The temperature anisotropy power spectrum are sums of tensor and scalar contributions, $C_l^{TT} = C_l^{TT,scalar} + C_l^{TT,tensor}$. To establish the amplitude of tensor fluctuations from total temperature anisotropies, which is what we observe, one must need to properly account for the contribution from scalar fluctuations. Thus, an independent estimate of the scalar amplitude, when combined with temperature anisotropy spectrum, can be used to constrain the amplitude of tensor fluctuations, or, the ratio of tensor-to-scalar fluctuation amplitude. We should always keep in mind that due to the integrated Sachs-Wolfe effect produced at late time by the scalar perturbations, $C_l^{TT,scalar}$ has a non-trivial normalization that depends both on the potential at last scattering (the Sachs-Wolfe effect), as well as the time evolution of the potential; The normalization of the CMB temperature power spectrum also depends on the amount as well as the nature of the dark energy given that potential fluctuations evolve differently depending on late-time cosmology [14, 15, 16].

The scalar amplitude can be inferred from the matter power spectrum as measured by galaxy surveys [17, 18]. Alternatively we can make use of the positive cross-correlation between WMAP anisotropy maps and LSS surveys [19, 20, 21, 22, 23]. Here we consider such a possibility and determine an independent constrain on the tensor-to-scalar ratio. Our limit improves those inferred from the combination of the WMAP data with either 2dF or SDSS surveys [10, 17], by nearly a factor of two. In particular we find that in a flat cosmological constant dominated universe $r < 0.59$ at 2σ . We also discuss how the constraints on r change by including more accurate information on the ISW effect.

II. LIMIT ON TENSOR-TO-SCALAR RATIO

The correlation between the large angular scale CMB temperature anisotropy maps and the distribution of the large scale structure (LSS) is consequence of the integrated Sachs-Wolfe. As the universe starts deviating from the matter dominated expansion, the gravitational potentials associated with clumps of matter change in time. CMB photons crossing these regions undergo a shift which causes a temperature anisotropy. Since the clustered structures arises from the scalar fluctuations the CMB-LSS correlation depends only on the scalar modes and does not get any contribution from the tem-

perature anisotropies induced by the gravitational waves background. Indeed the latter is uncorrelated with the distribution of galaxies, cluster and super-cluster. Thus, the cross-correlation between CMB maps and surveys can be used to infer A_s , under the assumption of a given dark energy model.

The angular cross-correlation in Legendre series is given by

$$C^X(\theta) = \sum_{l=2}^{\infty} \frac{2l+1}{4\pi} C_l^X P_l(\cos(\theta)), \quad (1)$$

where $P_l(\cos\theta)$ are the Legendre polynomials and C_l^X is the cross-correlation power spectrum [24, 26]:

$$C_l^X = \frac{2}{\pi} \int k^2 dk P_{\delta\delta}(k) I_l^{ISW}(k) I_l^{LSS}(k). \quad (2)$$

Here, $P_{\delta\delta}(k)$ is the power spectrum of density fluctuations (which captures the amplitude of scalar fluctuations) and integrand functions $I_l^{ISW}(k)$ and $I_l^{LSS}(k)$ are defined as

$$I_l^{ISW}(k) = \int W^{ISW}(k, r) j_l[kr] dr \quad (3)$$

$$I_l^{LSS}(k) = \int W^{LSS}(k, r) j_l[kr] dr \quad (4)$$

when the two window functions are

$$\begin{aligned} W^{ISW}(k, r) &= -3\Omega_m \left(\frac{H_0}{k}\right)^2 \frac{d}{dr} \left(\frac{G}{a}\right) \\ W^{LSS}(k, r) &= b(k, r) n(r) G(r). \end{aligned} \quad (5)$$

In above, $j_l[kr]$ is the spherical Bessel function when $r(z)$ is the comoving distance, $b(k, r)$ is the scale- and redshift, or distance-dependent bias factor of galaxies with a normalized distribution, in distance, of $n(r)$ ($\int dr n(r) = 1$), and $G(r)$ is the growth function of density fluctuations. Other parameters have standard definitions. In converting time evolving potential fluctuations, captured by the ISW effect, to density fluctuations, we have assume a flat-cosmological model. If not, the window function related to the ISW term, $W^{ISW}(k, r)$, will contain corrections related to curvature. We refer the reader to Ref. [24, 25, 26, 27, 28] for a more detailed discussion on the cross-correlation power spectrum.

From Eq. (4), it is evident that the determination of A_s from cross-correlation measurements depends on the galaxy bias $b(k, z)$ of the LSS catalog used. Nevertheless, we can state in a conservative way that the r.m.s. value of the matter fluctuation at $R = 8\text{Mpc}/h$ lies in the range $\sigma_8 = 0.7 - 1.1$ [15]. Normalizing the predicted matter-matter power spectra to these values and comparing to the observed galaxy-galaxy correlation function one can determine the corresponding bias. Marginalizing over σ_8 introduces, for a single catalog, an extra $\sim 20\%$ error in the measurement of the ISW-correlation signal.




FIG. 1: Likelihood contours in the $\Omega_\Lambda - r$ plane from current ISW detections (see [33]). Flatness is assumed. The ISW signal is smaller for small values of the Cosmological Constant.

This error can be further reduced if one considers several ISW detections with catalogs affected by different biases, or, in the case of a single survey, if biasing can be determined as a function of the galaxy luminosity [29]. Here is worth remarking that the dependence of the cross-correlation on the bias is *linear*. Therefore, the error on the inference of A_s induced by the uncertainty on the bias is fractionally less here than in the standard method involving the combination of CMB anisotropy data and LSS clustering spectrum (e.g. [10]), as the full galaxy-galaxy correlation function *quadratically* depends on the galaxy bias factor.

Another caveat concerns the dark energy. Current measurements of the cross-correlation lack the accuracy to strongly constrain any dark energy parameters [27]. The next generation of large scale structure surveys will provide better constraints by measuring the correlation between the CMB and different redshift sample of galaxies [30]. As shown in [25, 27] the redshift behavior of the cross-correlation signal depends on the specific of the dark energy, while A_s is an overall normalization factor independent of the redshift.

The cosmological constant is the simplest dark energy model to best fit current data [10, 17, 18, 31, 32].

We therefore limit our analysis to flat LCDM cosmologies and use the current ISW detections as collected by [33]. These data consist of average angular cross-correlation measurements between WMAP temperature maps and 5 different LSS surveys. For a more detailed discussion on the evaluation of the likelihood statistics, the effect of systematics and the correlations between redshift bins we refer to [27]. We perform a likelihood analysis for the amplitude of the tensor modes A_T and marginalize over the following parameters: the cosmological constant in the range $0.6 < \Omega_\Lambda < 0.9$, the scalar

spectral index $0.8 < n_s < 1.3$ and a Hubble parameter $0.6 < h < 0.9$. We also vary σ_8 between 0.7 and 1.1. We assume a spatially-flat cosmological model, and matter density changes as Ω_Λ is varied with the condition that $\Omega_m = 1 - \Omega_\Lambda$. To obtain constraints on the tensor amplitude, we only make use of large scale, $10 < \ell < 40$, WMAP data. In order to match the region of angular scales probed by the ISW-correlation data and avoid biasing due to lack of power in WMAP quadrupole to the octopole, we have excluded the region of multipole $\ell < 10$. We also vary the baryon density, as a function of h , with a fixed value for $\Omega_b h^2$ of 0.02, but found little variation in our results to variations in Ω_b ; This is due to the fact that the analysis excludes acoustic peaks that are more sensitive to the exact baryon density, relative to matter, while at large scales, corresponding to the multipole range above, there is little or no sensitivity in the anisotropy spectrum to parameter Ω_b .

We plot the results in Fig. 1, which shows the 1 and 2σ contours in the $\Omega_\Lambda - r$ plane. As we may notice current data constrain the tensor contribution to $r < 0.59$ at 2σ confidence level. Since we have only 5 weak detections of the ISW-galaxy cross-correlation, which may be affected by unknown systematics, one should be careful in a direct comparison with other constraints in the literature. Nonetheless the results of this analysis already illustrates the potential of this method which captures all the information one can extract on the tensor amplitude given the knowledge of the galaxy distribution. In fact, the previous standard approach combines only the CMB and matter power spectrum. Similarly we use the galaxy power spectrum to constrain the bias parameter, but with the addition of the CMB-LSS cross-correlation. The only way to improve this method would be to consider the cross-correlation as a function of galaxy luminosity and redshift, since this would allow a better separation of the uncertainties related to bias and dark energy respectively.

As shown in Fig. 1, the constraint on the tensor-to-scalar ratio depends on Ω_Λ . In particular larger values of Ω_Λ allow a slightly greater tensor contribution. This trend is expected, in fact for low Λ values the ISW-correlation is smaller, therefore to match the observed cross-correlation larger values of the scalar amplitude are favoured and thus strongly limiting the tensor contribution when combined with the CMB normalization. On the contrary, increasing the dark energy density causes a larger ISW-correlation signal which will favour low values of A_s and consequently a larger contribution of the tensor modes. If future dark energy measurements, such as from Type Ia supernovae, are to point towards values of the cosmological constant around $\Omega_\Lambda \sim 0.5 - 0.6$, the current cross-correlation data already constrain the GW contribution to be less than 10% of the scalar amplitude. Beyond dark energy, we also found a minimal correlation of our constraint with the spectral index n_s over the range of values considered. Given the limited information from the cross-correlation we did not vary other parameters or included a running to the tilt.

It is clear that accounting for the CMB-LSS cross-correlation provide better limits on the tensor-to-scalar ratio than those derived from WMAP data alone, from which $r < 0.9$. However the improvement does not increase significantly as one includes further information on the CMB-LSS cross-correlation. To understand this point, consider the CMB temperature fluctuation at large angular scales,

$$\Delta T = \Delta T^{\text{SW}} + \Delta T^{\text{ISW}} + \Delta T^{\text{GW}}. \quad (6)$$

Assuming that all other cosmological parameters are known except for the amplitude of the tensor perturbation A_T , the limit on the tensor-to-scalar ratio is

$$\sigma_{A_T}^{-2} = \sum_l \frac{1}{\sigma_{C_l}^2} \left(\frac{\partial C_l^{\text{GW}}}{\partial A_T} \right)^2 \quad (7)$$

where

$$\sigma_{C_l} = \sqrt{\frac{2}{2l+1}} (C_l^{\text{SW}} + C_l^{\text{ISW}} + 2C_l^{\text{SW-ISW}}), \quad (8)$$

and we have assumed cosmic variance limited measurements. Taking the full anisotropy spectrum (which includes contributions beyond SW and ISW effects), the above estimate leads to a cosmic variance limit on the tensor-to-scalar ratio of 0.06 at the one-sigma confidence level. In order to improve this limit further, the CMB-LSS cross-correlation can be used to remove the ISW contribution. The extent to which the signal improves depends on how well C_l^{ISW} , and to a lesser extent the correlation between SW and ISW effects $C_l^{\text{SW-ISW}}$ can be removed from the data. If a fraction c of the ISW effect, in temperature maps, can be accounted for by the CMB-LSS cross-correlation, then the improved noise estimate for the calculation is

$$\sigma'_{C_l} = \sqrt{\frac{2}{2l+1}} [C_l^{\text{SW}} + (1-c)^2 C_l^{\text{ISW}} + 2(1-c)C_l^{\text{SW-ISW}}], \quad (9)$$

where we have ignored that the removal in the temperature maps also reduces the correlated part with ISW by a similar fraction though in reality, this may be different given the projection effects and the redshift dependence of the LSS maps used to clean the CMB maps from the ISW contribution. Assuming that one can safely remove a 50% contribution of the ISW effect, the improvement is at the level of 5%. Therefore removing the ISW allows us to push the tensor-to-scalar limit only to the level of 0.056.

While future constraints are not significant, the ISW removal allows us to include information related to other parameters, as the ultimate limit from temperature anisotropies requires perfect knowledge on all cosmological parameters other than the tensor amplitude. Thus, there are significant gains to be made with using the CMB-LSS correlations to improve the tensor-to-scalar limit. This is similar to the improvements one see when combining CMB data with LSS data. Note that, in future, best limits on the tensor-to-scalar ratio will come from CMB polarization observations, rather than temperature anisotropies. It is well-known that the direct signal in CMB B-models presents the best opportunity to establish the tensor contribution, even in the presence of a confusion with a lensing effect [34, 35, 36, 37, 38].

III. CONCLUSIONS

Large scale CMB anisotropies involve a combination of the scalar and tensor fluctuations, we have argued that the scalar amplitude can be independently determined through the CMB-galaxy cross-correlation, under the reasonable assumption that the large-scale clustering of galaxies is not affected by primordial gravitational waves. Previous limits on the tensor amplitude have considered only the matter power spectrum to determine the scalar amplitude. We have shown that an improved limit can be obtained by including the cross-correlation between CMB and large scale structure surveys, in addition to the galaxy power spectrum alone. Using recently measured cross-correlation data, we obtain a constraint $r < 0.5$ at 68 % confidence level on the tensor-to-scalar ratio. The data also allow us to exclude gravity waves at a level of a few percent, relative to the density field, in a low - Lambda dominated universe ($\Omega_\Lambda \sim 0.5$). This technique can account for the knowledge of additional cosmological parameters which are required to determined the amplitude of the tensor modes and helps converging on the ultimate tensor-to-scalar ratio limit of 0.05 from temperature anisotropies alone (assuming all other cosmological parameters are known).

Acknowledgments

We like to acknowledge Levon Pogorian and Enrique Gaztanaga for discussion and help. P.S.C. is supported by Columbia Academic Quality Fund. AM is supported by MURST through COFIN contract no. 2004027755.

-
- [1] M. S. Turner, Phys. Rev. D**55**, 435 (1997).
 - [2] S. Dodelson, W. H. Kinney and E.W. Kolb, Phys. Rev. D**56** 3207 (1997).

- [3] L. M. Krauss, M. White, Phys. Rev. Lett. **69**, 869 (1992).
- [4] M. B. Hoffman, M. S. Turner, Phys. Rev. D**64**, 023506 (2001), astro-ph/0006312.

- [5] J. R. Pritchard and M. Kamionkowski, astro-ph/0412581.
- [6] H. V. Peiris *et al.*, Astrophys. J. Suppl. **148**, 213 (2003), astro-ph/0302225.
- [7] W. H. Kinney, E. W. Kolb, A. Melchiorri and A. Riotto, Phys. Rev. D **69**, 103516 (2004), hep-ph/0305130.
- [8] V. Barger, H. S. Lee and D. Marfatia, Phys. Lett. B **565**, 33 (2003), hep-ph/0302150.
- [9] S. M. Leach and A. R. Liddle, Phys. Rev. D **68**, 123508 (2003), astro-ph/0306305.
- [10] D.N. Spergel *et al.*, Astrophys. J. Supp. **148**, 175 (2003).
- [11] U. Seljak *et al.*, arXiv:astro-ph/0407372.
- [12] M. Kamionkowski and A. Kosowsky, Phys. Rev. D **57**, 685 (1998).
- [13] T. L. Smith, M. Kamionkowski and A. Cooray, arXiv:astro-ph/0506422; C. Ungarelli, P. Corasaniti, R. A. Mercer and A. Vecchio, arXiv:astro-ph/0504294; A. Cooray, arXiv:astro-ph/0503118.
- [14] P.S. Corasaniti, B.A. Bassett, C. Ungarelli and E.J. Copeland, Phys. Rev. Lett. **90**, 091303 (2003).
- [15] M. Kunz, P.S. Corasaniti, D. Parkinson and E. Copeland, Phys. Rev. D **70**, 041301 (2004).
- [16] A. Cooray, D. Huterer and D. Baumann, Phys. Rev. D **69**, 027301 (2004), astro-ph/0304268.
- [17] M. Tegmark *et al.* [SDSS Collaboration], Phys. Rev. D **69**, 103501 (2004), astro-ph/0310723.
- [18] U. Seljak *et al.*, arXiv:astro-ph/0407372.
- [19] P. Fosalba, E. Gaztanaga and F.J. Castander, Astrophys. J. **89**, L597 (2003).
- [20] S. Boughn and R. Crittenden, astro-ph/0305001; S. Boughn and R. Crittenden, Nature **427**, 45 (2004).
- [21] R. Scranton *et al.*, astro-ph/0307335.
- [22] M.R. Nolta *et al.*, Astrophys. J. **608**, 10 (2004).
- [23] N. Afshordi, Y. Loh and M.A. Strauss, Phys. Rev. D **69**, 083542 (2004).
- [24] J. Garriga, L. Pogosian and T. Vachaspati, Phys. Rev. D **69**, 063511 (2004).
- [25] L. Pogosian, astro-ph/0409059.
- [26] A. Cooray, Phys. Rev. D **65**, 103510 (2002), astro-ph/0112408.
- [27] P. S. Corasaniti, T. Giannantonio and A. Melchiorri, astro-ph/0504115.
- [28] N. Afshordi, Phys. Rev. D **70**, 083536 (2004), astro-ph/0401166.
- [29] I. Zehavi *et al.* [The SDSS Collaboration], arXiv:astro-ph/0408569.
- [30] L. Pogosian, P.S. Corasaniti, C. Stephan-Otto, R. Crittenden and R. Nichol, in preparation.
- [31] P.S. Corasaniti, M. Kunz, D. Parkinson, E.J. Copeland and B.A. Bassett, Phys. Rev. D **70**, 083006 (2004).
- [32] A.G. Riess *et al.*, Astrophys. J. **607**, 665 (2004); R. A. Knop *et al.*, Astrophys. J. **598**, 102 (2003), astro-ph/0309368; Y. Wang and M. Tegmark, Phys. Rev. Lett. **92**, 241302 (2004), astro-ph/0403292; D. Huterer and A. Cooray, Phys. Rev. D **71**, 023506 (2005), astro-ph/0404062.
- [33] E. Gaztanaga, M. Manera, T. Multamaki, astro-ph/0407022.
- [34] M. Kesden, A. Cooray and M. Kamionkowski, Phys. Rev. Lett. **89**, 011304 (2002), astro-ph/0202434.
- [35] L. Knox and Y. S. Song, Phys. Rev. Lett. **89**, 011303 (2002), astro-ph/0202286.
- [36] W. Hu and T. Okamoto, Astrophys. J. **574**, 566 (2002), astro-ph/0111606.
- [37] A. Cooray and M. Kesden, New Astron. **8**, 231 (2003), astro-ph/0204068; M. H. Kesden, A. Cooray and M. Kamionkowski, Phys. Rev. D **67**, 123507 (2003), astro-ph/0302536.
- [38] C. M. Hirata and U. Seljak, Phys. Rev. D **68**, 083002 (2003), astro-ph/0306354; U. Seljak and C. M. Hirata, Phys. Rev. D **69**, 043005 (2004), astro-ph/0310163.
- [39] As in [6] we define r as the inflationary scalar/tensor ratio at $k_0 = 0.002 Mpc^{-1}$